

Corning® CoolCell® Products

Reference List

CORNING

There is an increasing trend in the use of primary and specialized cell types in 3D cell culture, cancer research, biobanking, and cell therapy applications. The sensitivity of these cell cultures necessitates standardized and reproducible cryopreservation techniques to ensure robust post-thaw cellular integrity and performance.

For example, stem cells require proper cryopreservation to optimize cell viability and ensure maintenance of the undifferentiated state¹. 3D cell cultures such as organoids can be cryopreserved intact, in fragments, or as dissociated cells at a recommended rate of -1°C/minute using a controlled rate freezer or freezing containers for optimal recovery². For cell therapy production, controlled freezing of the final drug product before long-term storage is essential to ensure consistent integrity and therapeutic effect³.

Corning CoolCell alcohol-free cell freezing containers ensure a standardized controlled rate of -1°C/minute cell freezing in a -80°C freezer, without alcohol or any fluids. The proprietary Corning CoolCell technology utilizes a thermo-conductive alloy core and highly-insulative outer material to precisely control the rate of heat removal and provide reproducible cell cryopreservation.

Corning CoolCell containers are a cost-effective and smaller-scale alternative to commonly used programmable freezers and demonstrate the same freezing accuracy⁴ for use in research to cell therapy clinical trials³. CoolCell cryogenic containers are proven for use with a variety of cell types including stem cells, primary cells, PBMC, cell lines, organoids, and encapsulated cells.

The selection below highlights some of the recent publications in which Corning CoolCell containers were used. A list of additional citations follows. For more information about Corning CoolCell products visit www.corning.com/lifesciences.

1. Cohen R, et al. Standardized cryopreservation of pluripotent stem cells. *Current Protocols in Stem Cell Biology* 2014;1C.14.1-1C.14.14. doi: 10.1002/9780470151808.sc01c14s28.
2. Clinton J, and McWilliams-Koeppen, P. Initiation, expansion, and cryopreservation of human primary tissue-derived normal and diseased organoids in embedded three-dimensional culture. *Current Protocols in Cell Biology* 2019;82:e66. doi: 10.1002/cpcb.66.
3. Foussat A, et al. Effective cryopreservation and recovery of human regulatory T cells. *Bioprocess Int* 2014;12(3)s:34-38.
4. Thompson M, et al. Validation of a novel portable freezing device in the optimal freezing of peripheral blood mononuclear cells for potential cell therapy use. *Cytotherapy* 2014;16(54):S29.



Application	Area	Cell/Organoid Type	Reference	
3D Cell Culture	Organoid Culture	Gastric (human and mouse)	Bartfield S, et al. Organoids as model for infectious diseases: Culture of human and murine stomach organoids and microinjection of helicobacter pylori. <i>J Vis Exp</i> 2015;(105):e53359. doi: 10.3791/53359.	
		Human primary tissue-derived (various tissue types)	Clinton J and McWilliams-Koeppen P. Initiation, expansion, and cryopreservation of human primary tissue-derived normal and diseased organoids in embedded three-dimensional culture. <i>Curr Protoc Cell Biol</i> 2019;82(1):e66. doi: 10.1002/cpcb.66.	
		Intestinal crypt-derived (various animals)	Powell R, et al. WRN conditioned media is sufficient for in vitro propagation of intestinal organoids from large farm and small companion animals. <i>Biol Open</i> 2017;6:698-705. doi: 10.1242/bio.021717.	
		Liver and pancreas (mouse adult)	Broutier L, et al. Culture and establishment of self-renewing human and mouse adult liver and pancreas 3D organoids and their genetic manipulation. <i>Nat Protoc</i> 2016;11:1724-1743. doi: 10.1038/nprot.2016.097.	
		LuCaP prostate cancer (patient-derived xenografts)	Beshiri M, et al. A PDX/organoid biobank of advanced prostate cancers captures genomic and phenotypic heterogeneity for disease modeling and therapeutic screening. <i>Clin Cancer Res</i> 2018;24(17):4332-4345. doi: 10.1158/1078-0432.CCR-18-0409.	
		Prostate normal and cancer tissue (human and mouse)	Drost J, et al. Organoid culture systems for prostate epithelial tissue and prostate cancer tissue. <i>Nat Protoc</i> 2016;11(2):347-358. doi: 10.1038/nprot.2016.006.	
Cell Therapy	Gastrointestinal	Biopsies from ileal and/or colonic tissue from patients with Crohn's disease	Wildenberg M, et al. Evaluation of the effect of storage condition on cell extraction and flow cytometric analysis from intestinal biopsies. <i>Journal of Immunological Methods</i> 2018;459:50-54.	
	Immune Cell	CD4+CD25- T Responder cells	Morales J, et al. Automated clinical grade expansion of regulatory T cells in a fully closed system. <i>Front Immunol</i> 2019;10:38. doi: 10.3389/fimmu.2019.00038.	
		Differentiated Natural Killer (NK) cells from cord blood CD34+ cells	Domogala A, et al. Cryopreservation has no effect on function of natural killer cells differentiated in vitro from umbilical cord blood CD34+ cells. <i>Cytotherapy</i> 2016;18:754-759. doi: 10.1016/j.jcyt.2016.02.008.	
		MNC fraction (human)	Harper H and Rich I. Measuring the potency of a stem cell therapeutic. In: Rich I (ed.) <i>Stem Cell Protoc. Methods Mol Biol</i> , vol 1235. Humana Press, New York, NY. doi: 10.1007/978-1-4939-1785-3_4, 33-48.	
		PBMC (Peripheral blood mononuclear cell)		Higdon L, et al. Virtual global transplant laboratory standard operating procedures for blood collection, PBMC isolation, and storage. <i>Transplant Direct</i> 2016;2:e101. doi: 10.1097/TXD.0000000000000613.
				Lemieux J, et al. A global look into human T cell subsets before and after cryopreservation using multiparametric flow cytometry and two-dimensional visualization analysis. <i>J Immunol Methods</i> 2016;434:73-82. doi: 10.1016/j.jim.2016.04.010.
				Tapia-Calle, et al. Distinctive responses in an in vitro human dendritic cell-based system upon stimulation with different influenza vaccine formulations. <i>Vaccines</i> 2017;5(3):21. doi: 10.3390/vaccines5030021
			Thompson M, et al. Validation of a novel portable freezing device in the optimal freezing of peripheral blood mononuclear cells for potential cell therapy use. <i>Cytotherapy</i> 2014;16(54):S29.	
		PBMC; Type 1 Tregs (Ag-Treg)	Foussat A, et al. Effective cryopreservation and recovery of human regulatory T cells. <i>Bioprocess Int</i> 2014;12(3)s:34-38.	
		Tumor biopsy-derived TIL	Poschke I, et al. Identification of a tumor-reactive T-cell repertoire in the immune infiltrate of patients with resectable pancreatic ductal adenocarcinoma. <i>Oncol Immunology</i> 2016;5(12):e1240859. doi: 10.1080/2162402X.2016.1240859.	
Tumor-infiltrating T cells (TIL) from tumor biopsies	Idorn M, et al. Transfection of tumor-infiltrating T cells with mRNA encoding CXCR2. In: Rhoads R (ed.) <i>Synthetic mRNA. Methods Mol Biol</i> , vol 1428. Humana Press, New York, NY. doi: 10.1007/978-1-4939-3625-0_17.			

Application	Area	Cell/Organoid Type	Reference
Cell Therapy	Neuronal	Primary cell cultures from Parkinson's disease biopsies	Xu H, et al. Neurotrophic factor expression in expandable cell populations from brain samples in living patients with Parkinson's disease. <i>FASEB J</i> 2015;27:4157-4168. doi: 10.1096/fj.12-226555.
	Other	Human amnion epithelial cells (hAECs)	Gramignoli R, et al. Isolation of human amnion epithelial cells according to current good manufacturing procedures. <i>Curr Protoc Stem Cell Biol</i> 2016;37:1E.10.1-1E.10.13. doi: 10.1002/cpsc.2.
		Murine heart endothelial cells (MHEC5-T)	von Bomhard A, et al. Cryopreservation of endothelial cells in various cryoprotective agents and media-Vitrification versus slow freezing methods. <i>PLoS ONE</i> 2016;11(2):e0149660. doi:10.1371/journal.pone.0149660.
		General	Buckler R, et al. Technological developments for small-scale downstream processing of cell therapies. <i>Cytotherapy</i> 2016;18:301-306.
	Pancreas	Native human pancreatic islets (intact or single cells)	Rawal S, et al. Long-term cryopreservation of reaggregated pancreatic islets resulting in successful transplantation in rats. <i>Cryobiology</i> 2017;76:41-50.
	Stem Cell	hESC and iPSC lines	Cohen R, et al. Standardized cryopreservation of pluripotent stem cells. <i>Curr Protoc Stem Cell Biol</i> 2014;28:1C.14.1-1C.14.14. doi: 10.1002/9780470151808.sc01c14s28.
		hPSC lines (various); hPSC-RPE and hPSC-LESC cells	Hongisto H, et al. Xeno- and feeder-free differentiation of human pluripotent stem cells to two distinct ocular epithelial cell types using simple modifications of one method. <i>Stem Cell Res Ther</i> 2017;8:291. doi: 10.1186/s13287-017-0738-4.
		mESC (germline-competent)	Czechanski A, et al. Derivation and characterization of mouse embryonic stem cells from permissive and nonpermissive strains. <i>Nat Protoc</i> 2014;9(3):559-574. doi: 10.1038/nprot.2014.030.
		MSC (clinical BM-derived)	Moll G, et al. Do cryopreserved mesenchymal stromal cells display impaired immunomodulatory and therapeutic properties? <i>Stem Cells</i> 2014;32(9): 2430-2442. doi: 10.1002/stem.1729.
		MSC (horse BM- and UCB-derived)	Bertoni L, et al. Intra-articular injection of 2 different dosages of autologous and allogeneic bone marrow- and umbilical cord-derived mesenchymal stem cells triggers a variable inflammatory response of the fetlock joint on 12 sound experimental horses. <i>Stem Cells Int</i> 2019;9431894. doi.org/10.1155/2019/9431894.
Various	MSC (human adipose-derived)	Hoogduijn M, et al. Effects of freeze-thawing and intravenous infusion on mesenchymal stromal cell gene expression. <i>Stem Cells Dev</i> 2016;25(8):586-597. doi: 10.1089/scd.2015.0329	
	D1-mesenchymal stem cells, murine C2C12 myoblasts, human insulin secreting cell line 1.1B4, human retinal pigment epithelial cell line Arpe-19 encapsulated in alginate microcapsules	Gurruchaga H, et al. Low molecular-weight hyaluronan as a cryoprotectant for the storage of microencapsulated cells. <i>Int J Pharm</i> 2018;548:206-216.	
Cancer	Breast cancer	Patient triple-negative breast cancer cells	Le Gallo M, et al. Tumor analysis: freeze-thawing cycle of triple-negative breast cancer cells alters tumor CD24/CD44 profiles and the percentage of tumor-infiltrating immune cells. <i>BMC Res Notes</i> 2018;11:401. https://doi.org/10.1186/s13104-018-3504-5 .
	Neuroblastoma	Primary neuroblastoma explants	Braekveld N, et al. Neuroblastoma patient-derived orthotopic xenografts reflect the microenvironmental hallmarks of aggressive patient tumours. <i>Cancer Lett</i> 2016;375:384-389.
	Osteosarcoma	MG63 cells encapsulated in alginate fibers	Cagol N, et al. Effect of cryopreservation on cell-laden hydrogels: Comparison of different cryoprotectants. <i>Tissue Eng</i> 2018; Part C:24(1):20-31. doi: 10.1089/ten.tec.2017.0258.

For more specific information on claims, visit the Certificates page at www.corning.com/lifesciences.

Warranty/Disclaimer: Unless otherwise specified, all products are for research use only. Not intended for use in diagnostic or therapeutic procedures. Corning Life Sciences makes no claims regarding the performance of these products for clinical or diagnostic applications.

For additional product or technical information, visit www.corning.com/lifesciences or call 800.492.1110. Outside the United States, call +1.978.442.2200 or contact your local Corning sales office.

CORNING

Corning Incorporated
Life Sciences

836 North St.
Building 300, Suite 3401
Tewksbury, MA 01876
t 800.492.1110
t 978.442.2200
f 978.442.2476

www.corning.com/lifesciences

ASIA/PACIFIC

Australia/New Zealand
t 61 427286832

Chinese Mainland
t 86 21 3338 4338
f 86 21 3338 4300

India
t 91 124 4604000
f 91 124 4604099

Japan

t 81 3-3586 1996
f 81 3-3586 1291

Korea

t 82 2-796-9500
f 82 2-796-9300

Singapore

t 65 6572-9740
f 65 6735-2913

Taiwan

t 886 2-2716-0338
f 886 2-2516-7500

EUROPE

CSEurope@corning.com

France

t 0800 916 882
f 0800 918 636

Germany

t 0800 101 1153
f 0800 101 2427

The Netherlands

t 020 655 79 28
f 020 659 76 73

United Kingdom

t 0800 376 8660
f 0800 279 1117

All Other European Countries

t +31 (0) 206 59 60 51
f +31 (0) 206 59 76 73

LATIN AMERICA

grupoLA@corning.com

Brazil

t 55 (11) 3089-7400

Mexico

t (52-81) 8158-8400